

**IV Iberian Congress on Biotechnology
I Ibero-American Meeting on Biotechnology**

BIOTEC'98

BOOK OF ABSTRACTS

Edited by
MANUEL MOTA and EUGÉNIO C. FERREIRA
Departamento de Engenharia Biológica, Universidade do Minho
4700 Braga, PORTUGAL



This volume contains abstracts presented at the IV Iberian Congress on Biotechnology, I Ibero-American Meeting on Biotechnology, from 12 to 15 July 1998 at the Universidade do Minho in Guimarães, Portugal.

The event was sponsored by:

British Council

EXPO'98

Fundação Luso-Americana para o Desenvolvimento

Fundação para a Ciência e Tecnologia

Governo Civil do Distrito de Braga

Instituto do Vinho do Porto

Sociedad Española de Biotecnología

Sociedade Portuguesa de Biotecnologia

Universidade do Minho

título: BIOTEC'98 - Book of Abstracts
edited by Manuel Mota and Eugénio C. Ferreira

edição: 1ª edição, Julho de 1998
ISBN: 972-97810-0-1
Depósito legal: 125387/98
Tiragem: 600 exemplares

publicado por: Departamento de Engenharia Biológica, Universidade do Minho
Campus de Gualtar, 4700 Braga, Portugal

© *Universidade do Minho*

Immobilised Particles in Gel Matrix-Type Porous Media. Homogeneous porous media model

Manuel Mota*, José A. Teixeira and Alexander Yelshin

Centro de Engenharia Biológica - IBQF, Universidade do Minho, 4700 Braga, Portugal

Keywords: gel, immobilised cells, diffusion, tortuosity, porosity, and simulation.

Diffusion in pure gels and gels with immobilised cells were analysed. A model of diffusion in immobilised cell gel assuming of homogeneous cells distribution in gel was improved by introducing a tortuosity value. By theoretical analysis and numerical modelling it was shown that tortuosity of a gel with immobilised cells is the result of two factors: 1) a tortuosity generated by cells, T_c , and 2) a tortuosity of the gel matrix, T_g . Both variables are function of cell volume fraction, ϕ_c . Hence, the total path length in the gel with immobilised cell is $L_\Sigma = T_c T_g L$ and total tortuosity will be $T_\Sigma = L_\Sigma / L = T_c \cdot T_g$.

Based on this approach it is possible to analyse the data for a gel + cells system, where in general tortuosity seems too large. It is shown in this work that the diffusivity $\eta = D_e / D_0$ for this system is a complex function of: 1) diffusivity in gel, η_g and 2) diffusivity in immobilised cells, η_c . Hence

$\eta_c = D_e / D_g$, or $D_e = \eta_c D_g$, and $D_e / D_0 = \eta_c \frac{D_g}{D_0} = \eta_c \eta_g$, where D_g is the diffusion coefficient

in pure gel matrix, and D_0 is the diffusion coefficient in bulk liquid. Diffusion in a pure gel without immobilised cells is represented by the equation $\eta_g = D_g / D_0 = \varepsilon_g / T_g$, where ε_g is the gel porosity, and T_g is the pure gel tortuosity (average ratio of molecule path to porous media thickness).

The total diffusivity, with tortuosity T_c and T_g depending on volume fraction of immobilised

cells $(1 - \phi_c) = \varepsilon_c$, is $\eta = \eta_g \eta_c = \frac{\varepsilon_g}{T_g} \cdot \frac{\varepsilon_c}{T_c(\varepsilon_c) \cdot T_g(\varepsilon_c)}$.

For the general case, the tortuosity created in the matrix by cells presence is $T_c(\varepsilon_c) = 1 / \varepsilon_c^\beta = 1 / (1 - \phi_c)^\beta$ and the tortuosity of gel matrix filled with cells is $T_g(\varepsilon) = 1 / \varepsilon_c^\gamma = 1 / (1 - \phi_c)^\gamma$. Thus: $D_e / D_0 = \eta_g \cdot (1 - \phi_c) \cdot (1 - \phi_c)^\beta (1 - \phi_c)^\gamma = \eta_g (1 - \phi_c)^\alpha$, and $\alpha = 1 + \beta + \gamma$, where β and γ are order values in the range from 0 to 1, and η_g is the ratio D_g / D_0 for pure gels. In the particular case when $T_g(\varepsilon_c) = T_c(\varepsilon_c) = T$, and $T \sim 1 / \sqrt{\varepsilon_c}$ the equation

becomes $\eta = \frac{D_e}{D_0} = \frac{\varepsilon_g}{T_g} \cdot \frac{\varepsilon_c}{T^2} = \eta_g \varepsilon_c^2$.

The developed model gave the possibility to describe the dependence of D_e / D_0 on ϕ_c . Comparison with numerous published experimental data showed a good approach. Not all experimental data could be fitted with this homogeneous model. These deviations case might be explained by non-homogeneous cell distributions inside the gel matrix.